

Differences between α -linolenic and linoleic acid supplementation on the redox status and cardiodynamic parameters of male and female *Wistar albino* rats

Kristina Radoman¹, Vladimir Živković², Tamara Nikolić³, Isidora Stojić³, Danijela Raičević⁴, Jovana Jeremić³, Ivan Srejšević² and Vladimir Jakovljević^{2,5,*}

¹ College of Health Studies, Podgorica, Montenegro

² University of Kragujevac, Serbia, Faculty of Medical Sciences, Department of Physiology, Kragujevac, Serbia

³ University of Kragujevac, Serbia, Faculty of Medical Sciences, Department of Pharmacy, Kragujevac, Serbia

⁴ University of Montenegro, Biotechnical Faculty, Department for Viticulture and Enology, Podgorica, Montenegro

⁵ IM Sechenov First Moscow State Medical University, Moscow, Russia

*Corresponding author: drvladakbg@yahoo.com

Received: August 10, 2017; Revised: September 24, 2017; Accepted: October 6, 2017; Published online: October 16, 2017

Abstract: The aim of present study was to investigate the difference between α -linolenic acid (ALA, omega-3) and linoleic acid (LA, n-6) on the redox status and cardiac function of the isolated rat heart. ALA or LA were administered for 6 weeks by gavage to all animals, which were randomly divided into 4 groups: male rats treated with a linoleic acid (M-LA), dose of 7.3 mg/kg/day; female rats treated with a linoleic acid (F-LA), dose of 7.3 mg/kg/day; male rats treated with an α -linolenic acid (M-ALA), dose of 165 mg/kg/day; female rats treated with α -linolenic acid (F-ALA), dose of 165 mg/kg/day. Using the Langendorff technique, markers of heart function were evaluated: the maximum and minimum rates of pressure development in the left ventricle (LV; dp/dt max, dp/dt min), systolic and diastolic left ventricle pressure (SLVP, DLVP, respectively), heart rate (HR) and coronary flow (CF). We measured the concentrations of prooxidative markers: nitrites (NO_2^-), superoxide anion radicals (O_2^-) and hydrogen peroxide (H_2O_2), as well as the index of lipid peroxidation (TBARS) in the plasma and effluent. In the lysate, we measured the concentrations of reduced glutathione (GSH), catalase (CAT) and superoxide dismutase (SOD). ALA more negatively influenced the isolated rat heart, especially in females. In contrast, the administration of LA was linked to more prominent oxidative stress, while the application of ALA was associated with improved activity of the antioxidative defense system (with better values in males).

Key words: α -linolenic acid; linoleic acid; redox status; cardiac contractility; rat

INTRODUCTION

Polyunsaturated fatty acids (PUFAs) are fatty acids (FAs) that contain more than one double bond in their carbon backbone [1]. Polyunsaturated FAs can be classified into different groups according to their chemical structure; essential FAs are all omega-3 and omega-6 methylene-interrupted fatty acids. Among omega-6 FA, linoleic acid (LA; 18:2) is the most common PUFA, whereas α -linolenic acid (ALA; 18:3), a *cis*-omega-3 PUFA, is the most prevalent omega-3 FA in the diet [2]. Alpha-linolenic acid (ALA), or *all-cis*-9,12,15-octadecatrienoic acid, is an 18-carbon omega-3 essential FA and is the precursor of both eicosapentaenoic acid (EPA) and docosahexaenoic

acid (DHA). Linoleic acid (LA), or *all-cis*-9,12-octadecadienoic acid, is an 18-carbon omega-6 essential FA used in the biosynthesis of arachidonic acid (AA) and, thus, some prostaglandins, leukotrienes (LTA, LTB, LTC), and thromboxanes (TXA) [3].

Almost forty years ago, it was first hypothesized that an increased dietary intake of omega-3 and omega-6 PUFAs from fish fat and plant oil could exert protective effects against several pathologies, including cardiovascular [4,5], neurodegenerative [6], neuropsychiatric [6] and inflammatory diseases [7], as well as against some cancer types [8]. Since then, a large number of preclinical studies have clarified molecular mechanisms underlying the preventive and therapeutic

effects of omega-3 PUFAs against atherosclerosis and cardiac arrhythmia [4-8], and cardiologists and other physicians have begun to prescribe them routinely [9].

There has been an enormous discontinuity in the intake of LA during the last century as the availability of vegetable oils containing LA has increased 20-fold, yet at the same time, the generally accepted benefit of LA in slowing the development of atherosclerosis has come into question [9-10]. While some studies showed that the highest category of LA intake corresponded to a 15% lower risk of coronary heart disease, others have observed no benefit of LA supplementation on coronary disease [9-12].

Since LA and ALA, as well as other PUFAs, are highly oxidizable molecules, there is concern about their use at high concentrations for the prevention of chronic diseases that have oxidative stress as an underlying mechanism [13]. Some *in vitro* and *in vivo* studies have demonstrated the protective effects of ALA and LA against oxidative stress and lipid peroxidation. However, only a few, conflicting studies have been conducted showing the antioxidant potential of these essential FAs [14-16]. Controversy still exists over the precise mechanisms by which essential fatty acids could modulate cardiovascular function.

Omega-3 PUFAs have been shown to upregulate the antioxidant system in various cells, counteracting oxidative stress as antiinflammatory and cardioprotective FAs, while the omega-6 PUFAs were shown to be proinflammatory [7,9,17]. The intake of omega 3, omega 6 and *trans*-FAs may influence testicular function and compromise cardiovascular function in males [17]. Data have suggested that the omega-3 PUFA status is inversely associated with type 2 diabetes in women but not in men, suggesting a sex-dependent effect of PUFAs [18]. Therefore, LA and ALA likely differentially modulate the redox status and cardiac function in different genders.

Using an *ex vivo* model of the isolated rat heart and biochemical analyses, we investigated the difference between the effects of ALA omega-3 and LA omega-6, on the redox status and cardiac function of rat heart. Additionally, this study also examined the controversy over the use of these FAs as preventive/curative tools for cardiovascular disease and pays special attention to the role of gender in this.

MATERIALS AND METHODS

Ethics statement

All research procedures were carried out in accordance with the European Directive for the Welfare of Laboratory Animals, No. 86/609/EEC and the principles of good laboratory practice (GLP). The protocol of the current study was approved by the Ethics Committee for experimental animal well-being of the Faculty of Medical Sciences at the University of Kragujevac, Serbia.

Animals and experimental design

This study was carried out using 48 adult male and female *Wistar albino* rats (24 weeks old; b.w. 550±50 g). Animals were housed under controlled environmental conditions, at 25°C and a 12-h light/dark cycle with *ad libitum* access to food and tap water. For 6 weeks, the FA ALA or LA were administered by gavage to all animals [19], which were randomly divided into 4 groups as follows: (i) male rats treated with linoleic acid (M-LA) at a dose of 7.3 mg/kg/day [20]; (ii) female rats treated with linoleic acid (F-LA) at a dose of 7.3 mg/kg/day; (iii) male rats treated with α -linolenic acid (M-ALA) at a dose of 165 mg/kg/day [21]; (iv) female rats treated with α -linolenic acid (F-ALA) at a dose of 165 mg/kg/day.

Isolated heart perfusion

On the 43rd day, after 6 weeks of treatment with the different FA, all animals were euthanized by cervical dislocation without the use of an anesthetic in order to prevent any drug interactions. The chest was then opened via midline thoracotomy. The hearts were immediately removed, immersed in cold saline and mounted on a stainless-steel cannula on the Langendorff perfusion apparatus that provided retrograde perfusion under gradually increasing coronary perfusion pressure (CPP from 40 to 120 cmH₂O). Krebs-Henseleit buffer was used for retrograde perfusion (in mmol/L: NaCl 118, KCl 4.7, CaCl₂·2H₂O 2.5, Mg-SO₄·7H₂O 1.7, NaHCO₃ 25, KH₂PO₄ 1.2, glucose 11 and pyruvate 2). The buffer was equilibrated with 95% O₂ and 5% CO₂, pH 7.4 at 37°C. Following initiation of the heart perfusion, the preparations were stabilized

within 30 min with a basal coronary perfusion pressure of 70 cmH₂O. Following the stabilization period, the perfusion pressure was reduced to 50 and then to 40 cmH₂O; the pressure was then gradually increased to 60, 80, 100 and 120 cmH₂O to establish coronary autoregulation. Testing started immediately after the control experiment to avoid unwanted, time-dependent consequences. With a sensor placed in the left ventricle, software was used to measure the cardio-dynamic markers at each perfusion pressure (40, 60, 80, 100 and 120 cmH₂O). Markers of heart function included maximum and minimum rates of pressure development in the LV (dp/dt max, dp/dt min), systolic and diastolic left ventricle pressure (SLVP, DLVP, respectively), heart rate (HR) and coronary flow (CF). We evaluated all of the mentioned parameters during hypoxic (40 cmH₂O), normoxic (60-80 cmH₂O) and hyperoxic conditions (100-120 cmH₂O) according to coronary perfusion pressure (CPP).

Biochemical analyses

Blood samples were collected after 42 days of PUFA treatment and as the rats were killed by exsanguination. Samples of the coronary venous effluent were collected after stabilization of the coronary flow and for each perfusion pressure.

Preparation of plasma and erythrocyte lysate

Blood was collected into vacutainer tubes with anticoagulant (sodium citrate) and centrifuged for 10 min at 2000 x g to separate the plasma. The obtained plasma was used for determination of prooxidative markers: nitrites (the amount of NO released), superoxide anion radicals (O₂⁻), and hydrogen peroxide (H₂O₂), for the indirect quantification of the index of lipid peroxidation with reactive thiobarbituric substances (TBARS), which were measured by spectrophotometric quantification in the plasma. These parameters were also measured in the effluent samples. The remaining packed red blood cells (RBC) were washed three times with normal saline to remove the buffy coat. Hemolysis was performed by pipetting out 1 mL of washed RBC suspension in ice cold distilled water. RBC ghosts were sedimented in a high-speed refrigerated centrifuge at 10000 x g for 40 min. The cell

content was separated carefully and used for reduced glutathione (GSH), catalase (CAT) and superoxide dismutase (SOD) estimation. For hemoglobin (Hb) estimation we used the lysate and an equal volume of Drabkin's reagent, which were mixed and left for 3 min to determine the hemoglobin concentration. Absorbance was read at 540 nm against a reagent blank. Hemolysates containing about 50 g Hb/L and CAT, SOD and GSH were calculated as U/Hbx10³ according to McCord and Fridovich [22].

Determination of prooxidative parameters: superoxide anion radicals (O₂⁻), nitrites (NO₂⁻), hydrogen peroxide (H₂O₂), and the index of lipid peroxidation measured as TBARS

The concentration of the O₂⁻ radical was measured using the nitro blue tetrazolium (NBT) reagent in TRIS buffer (assay mixture) with coronary venous perfusate or plasma. The measurement was performed at 530 nm. The Krebs-Henseleit solvent served as the blank for effluent samples, and distilled water was the blank for plasma samples [23].

Nitric oxide (NO) rapidly decomposes into stable nitrite and nitrate metabolites. Nitrites can therefore be used as an index of NO production via a spectrophotometric method that uses the Griess reagent. Briefly, 0.5 mL of the perfusate or plasma was precipitated with 200 µL of 30% sulfosalicylic acid, mixed for 30 min and centrifuged at 30000 x g. Equal volumes of the supernatant and Griess reagent were mixed and stabilized for 10 min in the dark and the sample was measured spectrophotometrically at 543 nm. Nitrite concentrations were determined using sodium nitrite as the standard [24].

Hydrogen peroxide (H₂O₂) measurement was based on the oxidation of phenol red by H₂O₂ in a reaction catalyzed by horseradish peroxidase (HRP). A total of 200 µL of perfusate or plasma was precipitated with 800 µL of freshly prepared phenol red solution, followed by the addition of 10 µL of (1:20) HRP (made *ex tempore*). For the blank, Krebs-Henseleit solution or distilled water was used. The concentration of H₂O₂ was measured at 610 nm [25].

The index of lipid peroxidation was determined indirectly by measuring the products of the reaction

with thiobarbituric acid (TBA) or reactive substances (RS). Briefly, 1% TBA in 0.05 M NaOH was incubated with coronary venous perfusate or plasma at 100°C for 15 min and then spectrophotometrically measured at 530 nm. The Krebs-Henseleit solvent or distilled water served as the blanks [26].

Determination of antioxidant parameters

In the RBC lysate, we determined the concentrations of nonenzymatic antioxidants, such as reduced glutathione (GSH), and the activities of the antioxidant enzymes, catalase (CAT) and superoxide dismutase (SOD).

CAT activity was determined according to Beutler [27]. The lysates were diluted with distilled water (1:7 v/v) and treated with chloroform-ethanol (0.6:1 v/v) to remove Hb. Fifty μL of CAT buffer, 100 μL of sample and 1 mL of 10 mM H_2O_2 were mixed. Detection was performed at 360 nm.

SOD activity was determined by the McCord method [22]. The lysate (100 μL) and 1 mL of carbonate buffer were mixed together and 100 μL of epinephrine was added. Detection was performed at 470 nm.

The concentration of reduced glutathione (GSH) was determined based on GSH oxidation with 5,5'-dithiobis(2-nitrobenzoic acid), using the method by Beutler [28].

Drugs

α -Linolenic acid (product number L2376) and linoleic acid (product number L1376), both of $\geq 99\%$ purity, were purchased from Sigma-Aldrich Chemie GmbH, Eschenstr. 5, Taufkirchen, Germany.

Statistics

We used traditional parameters of descriptive statistics: average value \pm standard deviation (SD), and minimal and maximal values. Normality of the parameter distribution was evaluated with the Kolmogorov-Smirnov test. Dependent variables were compared using an analysis of variance (ANOVA) for repeated measurements (*post hoc* Bonferroni test). For independent groups, we used ANOVA in one direction (*post hoc* Tukey test). Statistical significance was based

on $p < 0.05$. Complete statistical evaluation was performed with SPSS Statistics 22 (SPSS, Chicago, IL).

RESULTS

Differences between chronic treatments with LA and ALA on the cardiodynamic parameters of isolated hearts of male and female rats

The values of dp/dt min (at all CPPs) and SLVP (60 cm CPP) after chronic administration of LA were significantly lower in male than in female rats (Fig. 1A-D). The values of CF were higher in the male group when compared with the female group under hypoxic (40-60 cm CPP) and normoxic conditions (80 cm CPP) (Fig. 1F). There were no differences in HR between the groups (Fig. 1E). Females treated with ALA had lower values of dp/dt max than males only under hypoxic conditions (Fig. 1A). In addition, CF and HR were also lower at all CPPs (Fig. 1 E and F).

In males, the values of SLVP (40, 60 cm CPP) and DLVP (at all CPPs) were lower in the ALA group as compared to the LA group (Fig. 1C and D). In contrast, ALA enhanced the HR and CF only in hypoxic conditions as compared with LA (Fig. 1E and F). In females, the values of all examined cardiodynamic parameters (except HR) were lower in the ALA group when compared to the LA group at all CPPs (Fig. 1A-D). CF was also lower under hyperoxic conditions (100 and 120 cm CPP) (Fig. 1E and F).

Differences between chronic treatments with LA and ALA on the biomarkers of oxidative stress in isolated heart (male and female rats)

Male rats treated with LA had higher values of H_2O_2 and NO_2^- (in hypoxic conditions) when compared to female rats (Fig. 2 B and D). In contrast, under normoxic and hyperoxic conditions, the females had higher values of O_2^- and TBARS (Fig. 2A and C). Male rats treated with ALA showed a larger release of all examined biomarkers of oxidative stress (Fig. 2A-D).

Male groups treated with ALA had higher values of measured cardiac oxidative markers as compared to the LA groups (Fig. 2A-D). The values of NO_2^- were higher only in hypoxic conditions (Fig. 2D), and of




Fig.1. The effects of chronic administration of LA and ALA on the cardio-dynamic parameters of isolated rat heart. **A** – The maximum rate of pressure development in the left ventricle (dp/dt max); **B** – the minimum rate of pressure development in the left ventricle (dp/dt min); **C** – SLVP; **D** – DLVP; **E** – HR; **F** – CF. All values are expressed as the mean \pm standard deviation. Statistically significant differences ($p < 0.05$) between the groups at the same coronary perfusion pressure are marked as follows: *a* – differences between the sexes treated with LA, *b* – differences between the sexes treated with ALA, *c* – differences between male rats treated with LA or ALA, *d* – differences between female rats treated with LA or ALA.




Fig. 2. The effects of the chronic administration of LA and ALA on the biomarkers of oxidative stress in isolated rat heart. **A** – O_2^- concentration; **B** – H_2O_2 concentration; **C** – the index of lipid peroxidation, measured as TBARS; **D** – the concentration of nitrites (NO_2^-).

O_2^- in normoxic and hyperoxic conditions (Fig. 2A). Female groups treated with ALA exhibited a stronger release of H_2O_2 (normoxic conditions) and TBARS (hypoxic and normoxic conditions) (Fig. 2B and C), and a lower release of O_2^- and NO_2^- (normoxic and hyperoxic conditions) in comparison to the LA groups (Fig. 2A and D).

Differences between chronic treatments with LA and ALA on the biomarkers of oxidative stress in blood samples of male and female rats

Males treated with LA showed higher release of O_2^- and NO_2^- (Fig. 3A and D). There were no statistically significant differences in the production of H_2O_2 and TBARS between the sexes (Fig. 3B and C). Males treated with ALA also had higher values of O_2^- (Fig. 3A). There were no statistically significant differences in the production of other examined biomarkers between the sexes (Fig. 3B-D).

The male ALA group had lower values of TBARS and NO_2^- when compared to the LA group (Fig. 3C and D). There were no statistically significant differences in the production of O_2^- and H_2O_2 between the compared groups (Fig. 3A and B). In females, there were no statistically significant differences in the production of all of the examined biomarkers between groups (Fig. 3A-D).

Differences between chronic treatments with LA and ALA on the levels of antioxidant markers in blood samples of male and female rats

Males treated with LA had higher activities of SOD and CAT (Fig. 4A and B), while the concentration of GSH was lower (Fig. 4C). In contrast, males treated with ALA had a higher activity of SOD and concentration of GSH (Fig. 4A and C), whereas CAT activity was unchanged in both sexes (Fig. 4B).




Fig. 3. The effects of the chronic administration of LA and ALA on the levels of prooxidant markers in blood samples. **A** – O_2^- concentration; **B** – H_2O_2 concentration; **C** – the index of lipid peroxidation, measured as TBARS; **D** – the concentration of nitrites NO_2^- .




Fig. 4. The effects of the chronic administration of LA and ALA on the levels of antioxidant markers in blood samples, **A** – SOD; **B** – CAT; **C** – GSH.

The ALA group had higher values of SOD activity and GSH concentrations (Fig. 4A and C) and lower values of CAT activity when compared with the LA group (Fig. 4B). In females, there were no statistically significant differences in antioxidant markers between the compared groups (Fig. 4A-C).

DISCUSSION

The aim of the current study was to investigate the differences between the two FAs, α -linolenic acid (ALA, omega-3) and linoleic acid (LA, omega-6), on the redox status and cardiac function of rat heart. Additionally, this study aimed to evaluate the potential

gender-related differences in response to ALA and LA treatment.

The two major types of long-chain PUFAs, omega-3 and omega-6, were shown to have antagonistic effects [1-7]. While omega-3 PUFAs emerged as anti-inflammatory and cardioprotective, omega-6 PUFAs were shown to be proinflammatory. To test these assumptions and the differences between these FAs on cardiac contractility, we evaluated the changes in cardiac function after 6 weeks of FA treatment using the following markers of heart function: maximum and minimum rates of pressure development in the LV (dp/dt max, dp/dt min), SLVP, DLVP, HR and CF. Myocardial contractile dysfunction, a major manifestation of heart failure, can be revealed through the mechanical defects of left ventricular pressure development ($+dp/dt$ max) and the rate of relaxation, as measured by LV negative ($-dp/dt$ min).

The contractility parameters were lower in the F-ALA group, while the same parameters were not affected in male groups. The other important indicators used to assess LV (SLVP and DLVP) were higher in the F-LA group (compared with the F-ALA). HR had the most drastic decrease in the M-ALA group. Interestingly, CF was lower in male animals treated with LA in contrast to female rats. When comparing the effects of LA between the male and female rats, lusitropic force and cardiac perfusion was lower in the female group. When comparing the effects of ALA between male and female rats, the only differences were observed in HR and CF.

A likely explanation for the cardiac effects of PUFA is that application of FA has profound effects on calcium management. PUFA can also produce a dose-dependent reduction in sodium and calcium currents by inhibiting the ryanodine receptor (RyR). These findings support the hypothesis that a lower omega-3 index may also be a marker of an increased propensity for the hypertensive rat heart towards malignant arrhythmias [29]. Other studies have pre-

sented similar results [28]. In accordance with their results, our findings suggest that these two FAs induce changes in cardiac function of isolated rat heart and act in opposite manners between genders. Generally, compared with ALA, LA significantly affected cardiac function and strongly disturbed heart muscle function, especially in female animals. It is probable that the duration of FA treatment is one of the crucial factors in the development of cardiac dysfunction and consequently of heart hypertrophy.

LA intake may compromise the omega-3 PUFA status of the tissue since its conversion to omega-6 LC-PUFA shares a common enzymatic pathway with the n-3 family [30]. The ALA diet led to higher levels of omega-3 LC-PUFAs, including DHA in the brain and heart [30]. One investigator described the different distribution of omega-6 and omega-3 PUFAs throughout the whole rat body, and 18:2 omega-6 was the most concentrated in the heart (13 wt%); this may be one of the reasons for the diverse effects of LA and ALA on heart function in our study [31].

One more mechanism whereby LA and ALA can modulate cardiac function was demonstrated by Berecki et al [31]. They stated that membrane incorporation of PUFAs reduces the response to norepinephrine. In particular, incorporated PUFAs blunted the increase in the sarcoplasmic reticulum (SR) calcium content produced by norepinephrine and therefore attenuated the increase in transient calcium produced by norepinephrine.

To determine the mechanism by which these FAs act on cardiac function *in vitro*, we evaluated the effects of LA and ALA on the prooxidative markers and parameters of systemic redox balance in the heart. In our study, female animals treated with LA exhibited a higher release of O_2^- , while the male groups treated with ALA had higher values of H_2O_2 . On the other hand, TBARS levels were significantly changed in almost all groups, with a higher release of both TBARS and NO observed in all female groups after treatment with either LA or ALA. It is important to note that in the M-ALA group, we observed a higher rate of coronary flow and all the prooxidative parameters measured from the effluent had a positive correlation with coronary flow. Most likely, FA-induced cardiac dysfunction is a consequence of myocardial disorders

but not of coronary circulation pathologies or endothelial dysfunction.

In the third part of this study we decided to exclude the effects of coronary flow rate by measuring all prooxidative and antioxidant markers from rat blood. We found higher values of O_2^- after treatment with both FAs in the male group, and higher levels of TBARS and nitrites in the M-LA group. These results are in accordance with the results of other relevant studies. Di Nunzio et al. [16] investigated the effect of LA and ALA on oxidative stress and confirmed the assumption that LA strongly induces lipid peroxidation. It is known that omega-3 PUFAs are important components of cell membranes that affect their function, and an omega-3 deficiency is deleterious to health. They are, however, prone to lipid peroxidation due to their many double bonds [4-10]. The metabolic reactions within the omega-3 and omega-6 PUFA families take place in the cell endoplasmic reticulum, apart from the last reaction of β -oxidation, which takes place in peroxisomes, thereby requiring translocation of adequate substrates into this cell compartment. PUFA residues of membrane phospholipids are very sensitive to oxidation and the action of reactive oxygen species (ROS).

Omega-3 PUFAs are known to have antiinflammatory effects. Excess production of NO is associated with inflammation [4,5]. NO is synthesized from L-arginine by NO synthase (NOS) with NADPH and oxygen as cosubstrates [32]. Inducible NOS (iNOS) is the key enzyme that produces large amounts of NO from macrophages stimulated by the bacterial endotoxin lipopolysaccharide (LPS) and by proinflammatory cytokines such as interferon- γ (IFN- γ) and tumor necrosis factor (TNF) [33]. In the present study, NO was significantly higher after LA supplementation, but the elevated levels of this parameter were decreased after ALA treatment.

Furthermore, SOD activity and GSH were significantly higher in the ALA group, in contrast to CAT activity. Generally, all antioxidant markers were significantly higher in ALA male and female groups, while LA induced higher levels of antioxidant parameters in the male group. Although different omega-3 acids share common chemical characteristics, a previous work showed that EPA and DHA modulate the

activity of antioxidant enzymes in different manners and they have different impacts on cell oxidative status. Popovic et al. [34] suggested that CAT activity and nitrite concentrations in liver were significantly decreased after the EPA/DHA supplementation. On the other hand, sex-related differences in this investigation indicate that the cardiodepressive effects of LA were more obvious in female rats during hypoxic conditions. Generally, males had higher values of cardiac and systemic oxidative parameters which was accompanied by a compensatory higher mobility of antioxidative protection.

Differences in the effects of LA and ALA on male and female animals may be related to the characteristics of male and female organisms. First, a higher content of body fat is characterized by a favorable fatty acid composition [35]. Second, the intake of omega-3 PUFAs is positively related to testicular volume, while the intake of omega-6 PUFAs is inversely related to testicular volume.

Finally, both fatty acids achieved their effects in hypoxic and hyperoxic conditions, suggesting that the level of oxygenation can also be an important factor in the response of the heart to the applied PUFAs. Thus, it seems that during normoxic conditions the heart is more resistant to their supplementation. ALA more negatively influenced the isolated rat heart in females. The administration of LA was connected to more prominent oxidative stress, while the application of ALA was also associated with improved activity of the antioxidative defense system (displaying higher values in males).

CONCLUSION

The findings of present study may help to elucidate the difference between the effect of ALA and LA on the heart and on local and systemic oxidative stress. The obtained results demonstrate that these PUFAs had a cardiodepressive impact on the heart, especially ALA. ALA apparently possesses antioxidative properties. Moreover, the effects of both FA were gender-specific in terms of exhibiting more negative effects in females. Lastly, ALA can be considered a beneficial supplement in all conditions and pathophysiological processes characterized by increased ROS production.

Acknowledgments: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author contributions: KR and VJ designed the study, IS, DR, JJ and TN collected and analyzed all data, IS and VZ performed statistical analyses. All authors approved final version of manuscript.

Conflict of interest disclosure: All authors of the present paper confirm no actual or potential conflicts of interest, including any financial, personal, or other relationships with people or organizations.

REFERENCES

1. Spector A, Kim H. Discovery of essential fatty acids. *J Lipid Res.* 2015;56:11-21.
2. Das U. Biological significance of essential fatty acids. *J Assoc Physicians India.* 2006;54:309-19.
3. Mele M, Cannelli G, Carta G, Cordeddu L, Melis M, Murru E, Stanton C, Banni S. Metabolism of c9,t11-conjugated linoleic acid (CLA) in humans. *Prostaglandins Leukot Essent Fatty Acids.* 2013;89:115-9.
4. Lunn J, Theobald H. The health effects of dietary unsaturated fatty acids. *Nutr Bull.* 2006;31:178-224.
5. Simopoulos A. The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Exp Biol Med.* 2008;233:674-88.
6. Patterson E, Wall R, Fitzgerald G, Ross R, Stanton C. Health implications of high dietary omega-6 polyunsaturated fatty acids. *J Nutr Metab.* 2012;53:942-6.
7. Arsić A, Prekajski N, Vučić V, Tepšić J, Popović T, Vrvic M, Glibetić M. Milk in human nutrition: comparison of fatty acid profiles. *Acta Vet (Beogr).* 2009;59:569-78.
8. Du R, Zhong T, Zhang W, Song P, Song W, Zhao Y, Wang C, Tang Y, Zhang X, Zhang Q. Antitumor effect of iRGD-modified liposomes containing conjugated linoleic acid-paclitaxel (CLA-PTX) on B16-F10 melanoma. *Int J Nanomedicine.* 2014;9:3091-105.
9. Farvid M, Ding M, Pan A, Sun Q, Chiuve S, Steffen L, Willett W, Hu F. Dietary linoleic acid and risk of coronary heart disease: A systematic review and meta-analysis of prospective cohort studies. *Circulation.* 2014;130:1568-78.
10. Chowdhury R, Warnakula S, Kunutsor S, Crowe F, Ward H, Johnson L, Franco O, Butterworth A, Forouhi N, Thompson S, Khaw K, Mozaffarian D, Danesh J, Di Angelantonio E. Association of dietary, circulating, and supplement fatty acids with coronary risk: A systematic review and meta-analysis. *Ann Intern Med.* 2014;160:398-06.
11. Blasbalg T, Hibbeln J, Ramsden C, Majchrzak S, Rawlings R. Changes in consumption of omega-3 and omega-6 fatty acids in the United States during the 20th century. *Am J Clin Nutr.* 2011;93:950-62.
12. Ramsden C, Zamora D, Majchrzak-Hong S, Faurot K, Broste S, Frantz R, Davis J, Ringel A, Suchindran C, Hibbeln J. Re-evaluation of the traditional diet-heart hypothesis: Analysis of recovered data from Minnesota Coronary Experiment (1968-1973). *BMJ.* 2016;353:1246-8.

13. Miyashita K. Paradox of omega-3 PUFA oxidation. *Eur J Lipid Sci Technol.* 2014;116:1268-79.
14. Di Nunzio M, Valli V, Bordoni A. PUFA and oxidative stress. Differential modulation of the cell response by DHA. *Int J Food Sci Nutr.* 2016;67:834-43.
15. Espinosa-Diez C, Miguel V, Mennerich D, Kietzmann T, Sánchez-Pérez P, Cadenas S, Lamas S. Antioxidant responses and cellular adjustments to oxidative stress. *Redox Biol.* 2015;6:183-97.
16. Chinnadurai K, Kanwal K, Tyagi A, Stanton C, Ross P. High conjugated linoleic acid enriched ghee (clarified butter) increases the antioxidant and antiatherogenic potency in female Wistar rats. *Lipids Health Dis.* 2016;12:121-25.
17. Kukoba T, Shysh A, Moïbenko O, Kotsiuruba A, Kharchenko O. The effects of alpha-linolenic acid on the functioning of the isolated heart during acute myocardial ischemia/reperfusion. *Fiziol Zh.* 2006;52:12-20.
18. Hassan A, Ibrahim A, Mbodji K, Coëffier M, Ziegler F, Bounoure F, Chardigny M, Skiba M, Savoye G, Déchelotte P, Marion-Letellier R. An α -linolenic acid-rich formula reduces oxidative stress and inflammation by regulating NF- κ B in rats with TNBS-induced colitis. *J Nutr.* 2010;140:1714-21.
19. Sikorska-Wiśniewska M, Miła A, Śledziński T, Małgorzewicz S, Stepnowski P, Rutkowski B, Chmielewski M. Disorders of serum omega-3 fatty acid composition in dialyzed patients, and their associations with fat mass. *Ren Fail.* 2017;39:406-12.
20. Folino A, Sprio E, Di Scipio F, Berta N, Rastaldo R. Alpha-linolenic acid protects against cardiac injury and remodeling induced by beta-adrenergic overstimulation. *Food Funct.* 2015;6:2231-9.
21. Mitchell L, Grant F, Melchert B, Petty M, Kennedy R. Linoleic acid metabolites act to increase contractility in isolated rat heart. *Cardiovasc Toxicol.* 2002;2:219-30.
22. McCord M, Fridovich I. SOD enzyme function for erythrocyte. *J Biol Chem.* 1969;244:6049-55.
23. Auclair C, Voisin E. Nitroblue tetrazolium reduction. In: Greenwald RA, editor. *CRC handbook of methods for oxygen radical research.* CRC Press Boca Raton; 1985. p. 123-32.
24. Green L, Wagnr D, Glogowski J, Skipper P, Wishnok J, Tannenbaum S. Analysis of nitrate, nitrite and (15 N) nitrate in biological fluids. *Anal Biochem.* 1982;126:131-8.
25. Pick E, Keisari Y. A simple colorimetric method for the measurement of hydrogen peroxide produced by cells in culture. *J Immunol Methods.* 1980;38:161-70.
26. Ohkawa H, Ohishi N. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem.* 1979;95:351-8.
27. Beutler E. Catalase. In: Beutler E, editor. *Red cell metabolism: A manual of biochemical methods.* New York: Grune and Stratton; 1982. p. 105-6.
28. Beutler E, Duron O, Kelly B. Improved method for the determination of blood glutathione. *J Lab Clin Med.* 1963;61:882-8.
29. Bačová B, Seč P, Radošinská J, Certík M, Vachulová A, Tribulová N. Lower omega-3 index is a marker of increased propensity of hypertensive rat heart to malignant arrhythmias. *Physiol Res.* 2013;62:201-8.
30. Ristic-Medic D, Suzic S, Vucic V, Takic M, Tepsic J, Glibetic M. Serum and erythrocyte membrane phospholipids fatty acid composition in hyperlipidemia: effects of dietary intervention and combined diet and fibrate therapy. *Gen Physiol Biophys.* 2009;28:190-9.
31. Berecki G, Den Ruijter H, Verkerk A, Schumacher C, Baartscheer A, Bakker D. Dietary fish oil reduces the incidence of triggered arrhythmias in pig ventricular myocytes. *Heart Rhythm.* 2007;4:1452-60.
32. Blanchard H, Pédrone F, Boulier-Monthéan N, Catheline D, Rioux V, Legrand P. Comparative effects of well-balanced diets enriched in α -linolenic or linoleic acids on LC-PUFA metabolism in rat tissues. *Prostaglandins Leukot Essent Fatty Acids.* 2013;88:383-9.
33. Salem N, Lin Y, Moriguchi T, Lim S, Salem Jr N, Hibbeln J. Distribution of omega-6 and omega-3 polyunsaturated fatty acids in the whole rat body and 25 compartments. *Prostaglandins Leukot Essent Fatty Acids.* 2010;100:13-20.
34. Popović T, Borozan S, Arsić A, Martačić J, Vučić V, Trbović A, Mandić L, Glibetić M. Fish oil supplementation improved liver phospholipids fatty acid composition and parameters of oxidative stress in male Wistar rats. *J Anim Phys Anim Nutr.* 2012;96:1020-9.
35. Komatsu W, Ishihara K, Murata M, Saito H, Shinohara K. Docosahexaenoic acid suppresses nitric oxide production and inducible nitric oxide synthase expression in interferon-gamma plus lipopolysaccharide-stimulated murine macrophages by inhibiting the oxidative stress. *Free Radic Biol Med.* 2003;34:1006-16.