

Alpha lipoic acid treatment induces the antioxidant system and ameliorates lipid peroxidation in maize seedlings under osmotic stress

Rabiye Terzi¹, Neslihan Saruhan Güler^{2,*}, Funda Gül Güven¹ and Asim Kadioglu¹

¹ Department of Biology, Faculty of Science, Karadeniz Technical University, 61080, Trabzon, Turkey

² Department of Nutrition and Dietetics, Faculty of Health Sciences, Karadeniz Technical University, 61080, Trabzon, Turkey

*Corresponding author: neslihansaruhan@hotmail.com

Received: December 18, 2017; **Revised:** February 22, 2018; **Accepted:** March 20, 2018; **Published online:** April 4, 2018

Abstract: Plants are markedly affected by drought stress caused by fluctuations in global climate, reduction in rainfall and a decrease in soil fertility. Therefore, some mechanistic strategies to cope with adverse effects of drought stress are needed. Alpha lipoic acid (ALA), a potent antioxidant molecule, is known to function in abiotic stress tolerance. In the current study, we investigated the ALA-stimulated physiological role in tolerance to osmotic stress induced by polyethylene glycol in two maize (*Zea mays* L.) cultivars (cv. Helen and cv. Akpinar). Application of ALA increased the leaf water potential of maize cultivars under stressful and stress-free conditions but decreased lipid peroxidation and the hydrogen peroxide (H₂O₂) content. Additionally, enhanced activity of the antioxidant defense system was observed following ALA application. Exogenous ALA elevated the activities of enzymatic antioxidants such as superoxide dismutase (SOD), catalase (CAT), guaiacol peroxidase (GPX), glutathione reductase (GR) and monodehydroascorbate reductase (MDHAR) under osmotic stress as compared to seedlings not exposed to ALA. Conversely, ascorbate peroxidase (APX) activity was decreased by ALA application in both cultivars. Higher GR and MDHAR activities of both cultivars were simultaneously observed in ALA treatments under osmotic stress. Taken together, the data indicated that exogenous ALA may function in arranging resilience against osmotic stress by reducing oxidative damage through induction of the antioxidant machinery in maize cultivars.

Key words: alpha lipoic acid; antioxidant system; osmotic stress; tolerance; *Zea mays*

INTRODUCTION

Plants growing in the natural environment are often subjected to various abiotic stresses. Drought-affected land has increased more than two-fold in recent years, and it is considered as one of the biggest threats to agriculture in the near future. Basic agricultural practices are markedly affected by drought stress because of fluctuations in global climate, reduction in rainfall and a decrease in soil fertility. Therefore, several adaptations and mitigation strategies are required to cope with drought stress. The standard effect of drought stress is oxidative damage. Exposure to drought results in a disruption of balance between reactive oxygen species (ROS) production and removal or scavenging [1]. Overproduction of ROS is a prominent anomaly in plants under unfavorable conditions that can cause an oxidative burst, thus blocking plant growth and development, or even leading to death. Therefore, plants

always try to maintain a well-developed enzymatic and nonenzymatic antioxidant defense system that can confront the deleterious effects of ROS [2].

It is important to explore suitable mechanisms for developing drought-tolerant crops capable of producing a sufficient yield under adverse stress conditions. Several researchers have attempted to find methods to alleviate drought stress or overcome drought injury in plants. Among them, the exogenous application of substances such as osmoprotectants, phytohormones, antioxidants and trace elements is attracting considerable attention [3]. In this regard, one of the natural molecules known to prevent or retard oxidation is alpha lipoic acid (ALA). ALA (1,2-dithiolane-3-pentanoic acid, 1,2-dithiolane-3-valeric acid, or thiocetic acid) is a dithiol, which effectively provides protection against oxidative stress by virtue of its two sulfhydryl moieties. In addition, it is an antioxidant compound

present in all types of eukaryotic and prokaryotic cells. Antioxidant substances typically possess antioxidant properties in their reduced form. Among antioxidant molecules, ALA is unique because it can retain powerful antioxidant functions in both its reduced (ALA) and oxidized (dihydrolipoic acid, DHLA) forms. Furthermore, both ALA and DHLA have a metal-chelating capacity and can scavenge free radicals [4]. In addition, ALA plays a crucial role in energy metabolism. The research on ALA in plants has attracted considerable attention in recent years, and evidence indicates that ALA can act in plants similarly to the way it does in animals [5,6]. The presence of ALA is of particular importance under excess copper [6] and salinity [7]. Furthermore, it affects specific proteins and enhances salt-stress tolerance [8]. In addition, it may mitigate oxidative stress by modulating ion homeostasis and the antioxidant system in salt-stressed wheat seedlings [9].

Maize is of pivotal economic importance globally as it has basic characteristics that make it extremely desirable for human consumption, animal feedstuff, as well as biodiesel/bioethanol production to scale down the dependence on fossil fuels [10-12]. Maize production is, however, negatively affected by some environmental factors, in particular, by drought and salt stress. Given the fact that maize is an agroeconomically important crop worldwide, it plays a pivotal role in producing abiotic stress-resistant and high-quality maize plants in the current scenario of global climate fluctuations and population explosion.

Studies examining the effect of exogenous ALA on plants exposed to stress are few and the mechanisms of ALA in promoting stress tolerance in plants need to be elucidated. Although there are studies that evaluated the role of ALA in the recycling of other antioxidants in plants under stress [6,7], no studies have evaluated the response of exogenous ALA on the antioxidant system under osmotic stress conditions in plants. In this study, we hypothesized that ALA pretreatment can increase the tolerance of maize seedlings to osmotic stress induced by polyethylene glycol (PEG₆₀₀₀) by enhancing antioxidant activity and decreasing ROS levels in plants. Therefore, we investigated the influence of ALA as a signal molecule by examining whether exogenous ALA pretreatment can alleviate the adverse effects of osmotic stress and

its relations with antioxidant systems. Furthermore, we determined the comparative performance of two maize cultivars differing in their water status in order to understand their adaptive mechanisms under drought stress with or without ALA. To determine the relationship between ALA and the antioxidant system, osmotic stress was applied to detached leaves of maize cultivars for a short period. However, this study is the first to demonstrate the capability of ALA in protecting maize seedlings against osmotic stress and should contribute to the understanding of the role of ALA in promoting osmotic stress tolerance.

MATERIALS AND METHODS

Plant materials, growth, and alpha lipoic acid treatment

Two maize (*Zea mays* L.) cultivars, cv. Helen and cv. Akpinar, differing in water status, were obtained from Advanta Seed Production Company Istanbul, Turkey and the Black Sea Agricultural Research Institute, Eskisehir, Turkey, respectively. The seeds were surface sterilized with 0.1% HgCl₂ for 3 min, followed by repeated washings with sterilized distilled water. The seeds were then sown in plastic pots (14 cm height, 16 cm top and 11 cm bottom diameters) containing peat and sand (5:1). The seedlings were grown in a greenhouse (temperature 21°C±2; relative humidity 60%±5; light intensity 400 μmol m⁻² s⁻¹) for 25 days. The pots were watered with distilled water every second day. To obtain an effective ALA import to the seedlings, the seedlings were excised from parts of the upper soil and rinsed in distilled water in test tubes wrapped with aluminum foil for 1 h to mitigate the effects of wound stress because of excision [13]. The detached seedlings were divided into four groups for each maize cultivar and submerged in dilute Hoagland nutrition solution (1/10 dilution; pH 6.0) with or without 0.02 mM ALA for 8 h. They were then subjected to osmotic stress treatments or kept in the Hoagland nutrient solution. Osmotic stress was gradually applied by the addition of PEG₆₀₀₀ in three increasing doses at an interval of 12 h, until a water potential of -0.3 MPa was instated. After this, the maize seedlings were exposed to osmotic stress for another 24 h and were divided into four groups as follows: (i) CTRL – con-

trol seedlings only exposed to Hoagland solution; (ii) PEG – osmotic stress treatment with 10% PEG₆₀₀₀ with -0.3 MPa as the osmotic potential; (iii) ALA – exposed to Hoagland solution containing 0.02 mM ALA; (iv) ALA+PEG – osmotic stress combined with 0.02 mM ALA. After the treatments, the second leaves of the plants were used for assays.

Measurements of the water potential

Leaf water potential (Ψ_{leaf}) was measured with a C52 thermocouple psychrometer (Wescor, Inc., Logan, UT, USA).

Dry weight measurement

To obtain adequate data on the growth variations of the seedlings, the dry weight was measured. The leaves were harvested from four group pots and fresh weights of the leaves were determined. The samples were dried in an oven at 85°C for 72 h [14].

Lipid peroxidation

Lipid peroxidation was measured in terms of the malondialdehyde (MDA) content ($\epsilon=155 \text{ mM}^{-1} \text{ cm}^{-1}$), a product of lipid peroxidation, following the method of Heath and Packer [15].

Determination of the hydrogen peroxide content

The endogenous H_2O_2 content was determined according to the modified method of Velikova et al. [16]. Leaves (0.25 g) were ground in 3 mL of 5% trichloroacetic acid with 0.1 g of activated charcoal at 0°C. To 0.5-mL aliquots of the supernatant, 0.5 mL of 10 mM potassium phosphate buffer (pH 7.0) and 0.75 mL of 1 M KI were added. The absorbance was measured at 390 nm and the H_2O_2 content was expressed as $\mu\text{mol g}^{-1} \text{ FW}$.

Antioxidant enzyme assays

Frozen leaf samples (0.5 g) were ground to a fine powder in liquid N_2 . The powder was homogenized in 5 mL of 50 mM potassium phosphate buffer (pH 7.0) containing 1% polyvinylpyrrolidone and 1 mM EDTA. For the ascorbate peroxidase assay, 5 mM of ascorbic

acid were added to the buffer. The homogenate was centrifuged at 20000 $\times g$ for 20 min at 4°C, and the supernatant was used for enzyme assays.

SOD (EC 1.15.1.1) activity was measured (25°C) based on the method of Beauchamp and Fridovich [17]. GPX (EC 1.11.1.7) activity was measured according to the method of Urbanek et al. [18] at 470 nm (25°C, $\epsilon = 26.6 \text{ mM}^{-1} \text{ cm}^{-1}$). APX (EC 1.11.1.11) activity was determined according to the method of Nakano and Asada [19], which is based on ascorbate oxidation at 290 nm (25°C, $\epsilon=2.8 \text{ mM}^{-1} \text{ cm}^{-1}$). CAT (EC 1.11.1.6) activity was measured according to the method of Aebi [20] (25°C, $\epsilon=39.4 \text{ mM}^{-1} \text{ cm}^{-1}$) at 240 nm. GR (EC 1.6.4.2) activity was determined following the decrease in absorbance at 340 nm (25°C, $\epsilon=6.22 \text{ mM}^{-1} \text{ cm}^{-1}$) associated with NADPH oxidation [21]. MDHAR (EC 1.6.5.4) activity was determined by following NADH oxidation at 340 nm (25°C, $\epsilon=6.22 \text{ mM}^{-1} \text{ cm}^{-1}$) [22]. The protein content was measured (25°C) according to the method of Bradford [23]. Bovine serum albumin was used as a standard. Enzyme activities were expressed as units per milligram of protein.

Statistical analysis

All experiments were performed in triplicate with four biological replicates. Duncan's multiple range test was used to perform variance analysis of means by SPSS software for Microsoft Windows (Ver. 15.0, SPSS Inc., Chicago, IL, USA). Statistical significance of the means between control and treatments was evaluated at 5% ($P<0.05$) probability level.

RESULTS

The effect of ALA pretreatment on the leaf water status

The effects of ALA application on the water potential and dry weight of the maize cultivars are presented in Fig. 1. The leaf water potential (Ψ_{leaf}) of both cultivars decreased in the PEG seedlings in comparison to the control seedlings. Ψ_{leaf} values of cv. Helen and cv. Akpınar were decreased by 66% and 58% in the PEG seedlings as compared to the control seedlings,

respectively. However, ALA+PEG seedlings sustained the Ψ_{leaf} at a higher level than the PEG seedlings. ALA+PEG enhanced the values of Ψ_{leaf} in cv. Helen and cv. Akpinar by 14% and 11%, respectively, as compared to the PEG seedlings (Fig 1A). Ψ_{leaf} values of cv. Helen and cv. Akpinar were increased by 30% and 18% in the ALA seedlings as compared to the control, respectively (Fig 1A).

The dry weight of cv. Akpinar but not of cv. Helen increased after the ALA treatment in comparison with the control seedlings. PEG treatment resulted in lower dry weight in cv. Helen than that of the control, but the dry weight of cv. Akpinar did not change after the PEG treatment in comparison with the control (Fig. 1B). The dry weights of neither cultivar changed after the ALA pretreatment under stress as compared to seedlings that were not exposed to ALA.

Effects of ALA pretreatment on lipid peroxidation and H_2O_2 content

The exposure to osmotic stress greatly increased membrane damage of both maize cultivars. The MDA content increased in cv. Helen and cv. Akpinar, 1.63- and 1.30-fold, respectively. However, the exogenous application of ALA alleviated the membrane damage of both cultivars under osmotic stress. Membrane damage was reduced in cv. Helen and cv. Akpinar, 1.16- and 1.19-fold, respectively (Fig. 2A).

The osmotic stress caused up to 3.28-fold accumulation of H_2O_2 in cv. Helen and 2.63-fold in cv. Akpinar when compared to matching controls. This indicated that oxidative stress occurred in osmotically-stressed maize seedlings. Exogenous ALA alone reduced the increase in H_2O_2 accumulation in both cultivars. The ALA+PEG pretreatment decreased the H_2O_2 content when compared with the PEG treatment. The decrease caused by exogenous ALA was relatively more pronounced in cv. Helen than in cv. Akpinar (1.41- and 1.22-fold, respectively) (Fig. 2B).

Effects of ALA pretreatment on antioxidant enzymes

All antioxidant enzyme activities increased under PEG-induced osmotic stress in both cultivars, and this increase was higher in the ALA-pretreated seed-




Fig. 1. The effect of exogenous ALA on leaf water potential (Ψ_{leaf}) (A) and dry weight (B) in two maize cultivars, Helen (filled square) and, Akpinar (open square), under osmotic stress conditions. The seedlings were subjected to four different treatments: (i) the control seedlings were only exposed to Hoagland solution (CTRL); (ii) pretreated with ALA and not osmotically stressed (ALA); (iii) only osmotically-stressed (PEG); (iv) pretreated with ALA and osmotically stressed (ALA+PEG). Vertical bars represent standard deviation. The different letters denote significant differences between the two cultivars along with the different treatments ($P < 0.05$).




Fig. 2. The effects of ALA on lipid peroxidation (A) and hydrogen peroxide (H_2O_2) content (B) in two maize cultivars, Helen (filled square) and Akpinar (open square), under osmotic stress conditions. Vertical bars represent the standard deviation. The different letters denote significant differences between the two cultivars along with the different treatments ($P < 0.05$).




Fig. 3. The effects of ALA on the activities of antioxidant enzymes (SOD (A), CAT (B), GPX (C), APX (D), GR (E), MDHAR (F)) in two maize cultivars, Helen (filled square) and Akpinar (open square), under osmotic stress conditions. Vertical bars represent the standard deviation. The different letters denote significant differences between the two cultivars along with the different treatments at ($P < 0.05$).

lings. Exogenous PEG alone increased the SOD activity of both cultivars as compared with that of their controls. Increases in SOD activity by 14% and 11% were recorded in cv. Helen and cv. Akpinar under osmotic stress, respectively. However, the application of exogenous ALA alone did not cause a change in SOD activity. Furthermore, an increase in the activity was observed by the application of ALA+PEG when compared with that of PEG alone. Increases in SOD activity by 23% and 28% were recorded after ALA application in cv. Helen and cv. Akpinar, respectively, under osmotic stress (Fig. 3A).

Osmotic stress increased CAT activities in cv. Helen and cv. Akpinar; however, the effect was not as high as that of ALA treatment alone. The activity increased by 172% in PEG seedlings of cv. Helen, while it increased 14% in cv. Akpinar, as compared to the control, respectively. CAT activity was increased in ALA-treated seedlings compared with that in control seedlings in both cultivars. The ALA application

under osmotic stress considerably increased CAT activity in both cultivars when compared with the PEG-treated seedlings. Increases in activities by 6% and 27% were recorded in cv. Helen and cv. Akpinar, respectively (Fig. 3B).

The PEG treatment increased GPX activity in both cultivars. As compared to the control, the activity was increased by 26% in PEG-treated seedlings of cv. Helen while it increased 62% in cv. Akpinar. GPX activity of the greatest magnitude was observed in ALA+PEG-pretreated seedlings; GPX activity in ALA+PEG applied seedlings increased by 53% and 4% in cv. Helen and cv. Akpinar as compared to the PEG seedlings, respectively (Fig. 3C).

APX activity of both cultivars increased under osmotic stress. Increases in activity by 41% and 14% were recorded under osmotic stress in cv. Helen and cv. Akpinar, respectively. In contrast, the ALA treatment alone did not cause a change in APX activity of

cv. Helen; however, it led to a considerable decrease in APX activity in cv. Akpınar as compared with that of their controls. ALA+PEG treatment caused a decrease in APX activity of both cultivars as compared with the PEG treatment. In ALA+PEG-treated seedlings, the activity decreased by 12% and 30% in cv. Helen and cv. Akpınar, respectively (Fig. 3D).

GR activity of maize cultivars was induced by the PEG treatment. Increases in activity by 24% and 123% were recorded under osmotic stress in cv. Helen and cv. Akpınar, respectively. However, no considerable differences in GR activity in cv. Helen between control and ALA-treated seedlings were determined. GR activity of the greatest magnitude was observed after the ALA+PEG pretreatment. When compared to PEG-treated seedlings, increases in activities by 35% and 25% after ALA application were recorded in cv. Helen and cv. Akpınar exposed to osmotic stress, respectively (Fig. 3E).

In both maize cultivars, osmotic stress caused a decrease in MDHAR activities when compared to matching controls. In PEG-treated seedlings, the activity decreased by 11% and 19% in cv. Helen and cv. Akpınar, respectively. The application of exogenous ALA alone markedly reversed the reduction in MDHAR activity. The ALA+PEG treatment considerably increased MDHAR activity in both cultivars in comparison to the PEG treatment; however, the effect was not as high as that observed in seedlings treated only with ALA. In ALA+PEG-treated seedlings, the activity increased by 23% and 45% in cv. Helen and cv. Akpınar, respectively (Fig. 3F).

DISCUSSION

The protective mechanisms of ALA, an antioxidant molecule, in inducing osmotic stress tolerance in plants remain to be elucidated. The present investigation suggests that the exogenous application of ALA can help reduce the adverse effects of osmotic stress in maize cultivars. The water status of maize cultivars was reduced under osmotic stress, while the ALA pretreatment substantially enhanced the leaf water potential. The pretreatment with ALA further enhanced the leaf water potential in cv. Helen in comparison to cv. Akpınar. A strong correlation exists between the plant

water content and accumulation of compatible solutes under drought stress [24]. Therefore, the improvement in leaf water potential by the exogenous application of ALA may be the result of osmotic adjustment due to the accumulation of compatible solutes, such as proline and sugar. Indeed, our results show that the proline and total soluble sugar contents in ALA-treated plants were higher than in PEG-treated plants (unpublished data). In addition, our findings show that the improvement in water status following the exogenous application of ALA may be the result of the alleviation of the detrimental effects of osmotic stress. Similar to our findings, Görcek and Erdal [9] reported that exogenous ALA treatment mitigated the salt-induced decrease in water status and, hence, contributed by increasing the leaf surface area. In addition, the authors indicated that enhancement of the water status and leaf surface area due to foliar application of ALA was due to its modulating role in ion homeostasis, osmotic regulation and the antioxidant system. In this study, osmotic stress markedly reduced the dry weights of cv. Helen but not Akpınar. When treated with ALA alone, a slight increase in the dry weight of cv. Akpınar was observed as compared to the control. The applied ALA was easily taken up by maize seedlings because they were detached and did not have a root system. It could be transported to the leaves and thus exogenous ALA could increase the dry weight of the leaves during the experimental period. Similarly, Mohammadkhani and Heidari [25] reported that dry weight changed in maize cultivars exposed to PEG-induced water stress for 24 h. Also, no significant differences were observed in the dry weights of PEG- and ALA+PEG-treated seedlings in both cultivars. We can say that exogenous ALA may not affect dry weight of the leaves under short-term osmotic stress conditions.

Even under optimal conditions, several metabolic processes lead to ROS production. The production of toxic oxygen derivatives is increased in response to all types of abiotic or biotic stresses [26]. Under osmotic stress, the interaction between ALA and increased ROS levels remains unknown. The primary site of PEG-induced osmotic stress injury was probably the cell membrane. Our results indicate that osmotic stress induced substantial damage to cell membranes as a result of increased cellular levels of MDA and H_2O_2 in both cultivars. In addition, the H_2O_2 content

was highest in PEG-treated seedlings, indicating that the generation of endogenous H_2O_2 exceeded the capacity of the cellular antioxidant defense system to eliminate H_2O_2 [27]. However, the exogenous application of ALA considerably reduced the endogenous levels of H_2O_2 and MDA by inducing the antioxidant system. The reduction in H_2O_2 content after ALA application may be due to its ability to scavenge ROS, as antioxidant enzymes. This correlates with the findings of Gorcek and Erdal [9] who observed that ALA significantly reduced H_2O_2 and superoxide contents and contributed to the response to oxidative damage and metabolic distortions in salt-stressed wheat seedlings. Likewise, in another study, it was shown that ALA can have protective and antimutagenic effects against oxidants (particularly H_2O_2) in yeast cells via its antioxidant activity [28]. Indeed, using various model systems, ALA was found to be highly reactive against a variety of ROS *in vitro*. ALA at concentrations of 0.05-1 mM scavenged H_2O_2 , the hydroxyl radical, hypochlorous acid and singlet oxygen. ALA also formed stable complexes with Mn^{2+} , Cu^{2+} , Zn^{2+} and chelated Fe^{2+} . Several studies provided evidence that ALA application decreased oxidative stress and restored reduced levels of other antioxidants under different physiological conditions *in vivo* [4,29]. Furthermore, ALA is highly effective in protecting stress-induced lipid peroxidation. It has been reported that ALA acts as a free radical scavenger and plays a crucial role in the recycling of other oxidized radical scavengers such as glutathione and ascorbate [4,5]. The results suggest that the reduction in MDA content was because of increased antioxidant enzyme activities that reduced H_2O_2 levels and membrane damage.

Plants possess efficient systems for scavenging ROS that protect them from destructive oxidative reactions. As part of this system, antioxidant enzymes are key elements in defense mechanisms [26]. In this study, we found that osmotic stress increases antioxidant enzyme activities, except MDHAR, in both maize cultivars (Fig. 3). In addition, cv. Helen had higher APX and CAT activities and lower GPX and GR activities than cv. Akpınar. Moreover, the exogenous application of ALA under osmotic stress further increased the activities of all these enzymes in both cultivars, with higher increases in GPX and GR activities in cv. Helen than in cv. Akpınar (Fig. 3C and E). APX activ-

ity, which was highly increased under osmotic stress in cv. Helen, was decreased in ALA-treated seedlings under osmotic stress. Therefore, ALA application to the cv. Helen exposed to osmotic stress may not induce some antioxidant enzymes such as APX under stress. In addition, the decrease in APX activity may be a result of ALA using other antioxidant enzymes instead of APX to remove H_2O_2 . Furthermore, the inhibition of APX activity by ALA may be because of downregulated gene expression or degradation, denaturation, or inactivation of this protein. However, in ALA-pretreated seedlings, the higher activities of antioxidant enzymes (except APX) correspond to a decrease in the endogenous H_2O_2 level, suggesting that pretreatment with exogenous ALA increases the ability of maize cultivars to scavenge H_2O_2 through antioxidant enzymes under osmotic stress. Similarly, Görcek and Erdal [9] reported that ALA application resulted in further increases in enzyme activities when compared to salt-stressed wheat seedlings, suggesting that ALA may contribute to increased plant resistance against salinity by effectively scavenging ROS through enhanced antioxidant activity.

In this study, we observed a trend similar to that of enzymes for GR activity in both cultivars. However, MDHAR activity decreased in both cultivars under osmotic stress. GR and MDHAR activities of both cultivars were increased by ALA application under stress. The results of our experiment showed that GR and MDHAR played crucial roles in scavenging H_2O_2 under osmotic stress, although APX had been inactivated. In addition, these antioxidant activities in plants typically depend on antioxidant metabolites such as ascorbate, glutathione, tocopherol, and polyphenols [30]; in particular, ascorbate and glutathione play key roles in redox signal transduction in higher plants under stress conditions [22]. It has been reported that in addition to scavenging ROS, ALA strengthens the antioxidant network by recycling other antioxidants, such as ascorbate and reduced glutathione in both plants and animals [7]. Therefore, this metabolism may be partly responsible for the high antioxidant activity. In addition, higher GR and MDHAR activities of both cultivars were simultaneously observed after ALA treatments under osmotic stress, and an increase in GR activity can promote the recycling of GSSG to GSH [31]. It is well known that the func-

tion of ascorbate depends on its redox status, which is closely related to MDHAR activity, and the higher MDHAR activity induced by exogenous ALA application under osmotic stress is crucial for efficient antioxidant activity.

CONCLUSIONS

Maize is a valuable crop and it is very important to establish mechanisms for developing drought-tolerant plants that can produce sufficient yields under adverse stress conditions. In this study, we describe the protective roles of ALA in maize seedlings under osmotic stress via modulation of H₂O₂ accumulation, alleviation of the water status and reorganization of the antioxidant machinery. Exogenous ALA may be considered a potential signaling molecule for augmenting the antioxidant potential of maize plants under osmotic stress conditions. In our opinion, ALA may have high potential impact for crop quality and sustainable agriculture but we are just beginning to understand its role. In the coming years, we assume that ALA will be able to improve plant growth and yield quality by increasing the antioxidant capacity of plants under water stress, and could become a novel candidate for sustainable agriculture. In our future research, we plan to analyze transcript levels of genes involved in osmolytic metabolism by quantitative PCR.

Acknowledgments: This work was supported by the Research Unit of Karadeniz Technical University (BAP, 8922).

Author contributions: RT and FGG designed the study and performed the experiments. NSG and AK evaluated the data and wrote the paper.

Conflict of interest disclosure: The authors state that they have no conflict of interest to disclose in the subject matter or materials discussed in this manuscript.

REFERENCES

- Smirnov N. Plant resistance to environmental stress. *Curr Opin Biotechnol.* 1998;9:214-19.
- Hasanuzzaman M, Hossain MA, Teixeira da Silva JA, Fujita M. Plant responses and tolerance to abiotic oxidative stress: antioxidant defense is a key factors. In: Bandi V, Shanker AK, Shanker C, Mandapaka M, editors. *Crop stress and its management: perspectives and strategies.* Berlin, Germany: Springer; 2012. p. 261-316.
- Hasanuzzaman M, Alam M, Rahman A, Hasanuzzaman Md, Nahar K, Fujita M. Exogenous proline and glycine betaine mediated upregulation of antioxidant defense and glyoxalase systems provides better protection against salt-induced oxidative stress in two rice (*Oryza sativa* L.) varieties. *Biomed Res Int.* 2014;2014:1-17.
- Packer L, Witt EH, Tritschler HJ. Alpha-lipoic acid as a biological antioxidant. *Free Radic Biol Med.* 1995;19:227-50.
- Navari-Izzo F, Quartacci MF, Sgherri CLM. Lipoic acid: a unique antioxidant in the detoxification of activated oxygen species. *Plant Physiol Biochem.* 2001;40:463-70.
- Sgherri C, Quartacci MF, Izzo R, Navari-Izzo F. Relation between lipoic acid and cell redox status in wheat grown in excess copper. *Plant Physiol Biochem.* 2002;40:591-7.
- Perez-Lopez U, Robredo A, Lacuesta M, Sgherri C, Mena-Petite A, Navari-Izzo F, Munoz-Rueda A. Lipoic acid and redox status in barley plants subjected to salinity and elevated CO₂. *Physiol Plant.* 2010;139:256-68.
- Yildiz M, Akçali N, Terzi H. Proteomic and biochemical responses of canola (*Brassica napus* L.) exposed to salinity stress and exogenous lipoic acid. *J Plant Physiol.* 2015;179:90-9.
- Gorcek Z, Erdal S. Lipoic acid mitigates oxidative stress and recovers metabolic distortions in salt-stressed wheat seedlings by modulating ion homeostasis, the osmo-regulator level and antioxidant system. *J Sci Food Agric.* 2015;95:2811-17.
- Crutzen PJ, Mosier AR, Smith KA, Winiwarter W. N₂O release from agro-biofuel production negates global warming reduction by replacing fossil fuels. *Atmos Chem Phys.* 2008;8:389-95.
- Food and Agriculture Organization of the United Nations. *Save and grow in practice maize rice wheat. A guide to sustainable cereal production.* Rome: Food and Agriculture Organization of the United Nations; 2016. 109 p.
- Pehlivan N, Yesilyurt AM, Durmus N, Karaoglu SA. *Trichoderma lixii* ID11D seed biopriming mitigates dose dependent salt toxicity in maize. *Acta Physiol Plant.* 2017;39(3):79.
- Terzi R, Kalaycioglu E, Demiralay M, Saglam A, Kadioglu A. Exogenous ascorbic acid mitigates accumulation of abscisic acid, proline and polyamine under osmotic stress in maize leaves. *Acta Physiol Plant.* 2015;37:43.
- Fu J, Huang B. Involvement of antioxidants and lipid peroxidation in the adaptation of two cool-season grasses to localized drought stress. *Environ Exp Bot.* 2001;45:105-14.
- Heath RL, Packer L. Photoperoxidation in isolated chloroplast. I. kinetics and stoichiometry of fatty acid peroxidation. *Arch Biochem Biophys.* 1968;125:189-98.
- Velikova V, Yordanov I, Edreva A. Oxidative stress and some antioxidant systems in acid rain-treated bean plants, protective role of exogenous polyamines. *Plant Sci.* 2000;151:59-66.
- Beauchamp C, Fridovich I. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal Biochem.* 1971;44:276-87.
- Urbanek H, Kuzniak-Gebarowska E, Herka H. Elicitation of defense responses in bean leaves by *Botrytis cinerea* polygalacturonase. *Acta Physiol Plant.* 1991;13:43-50.
- Nakano Y, Asada K. Hydrogen peroxide is by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.* 1981;22:867-80.

20. Aebi H. Catalase. In: Bergmeyer H, editor. Methods of enzymatic analysis. Chemie: Weinheim-Verlag; 1983. p. 273-86.
21. Foyer CH, Halliwell B. Presence of glutathione and glutathione reductase in chloroplast: a proposed role in ascorbic acid metabolism. *Planta*. 1976;133:21-5.
22. Hossain MA, Nakano Y, Asada K. Monodehydroascorbate reductase in spinach chloroplasts and its participation in regeneration of ascorbate for scavenging hydrogen peroxide. *Plant Cell Physiol*. 1984;25:385-95.
23. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities protein utilizing the principle of protein-dye binding. *Anal Biochem*. 1976;72:248-54.
24. Farooq M, Basra SMA, Wahid A, Ahmad N, Saleem BA. Improving the drought tolerance in rice (*Oryza sativa* L.) by exogenous application of salicylic acid. *J Agron Crop Sci*. 2009;195:237-46.
25. Mohammadkhani N, Heidari N. Water stress induced by polyethylene glycol 6000 and sodium chloride in two maize cultivars. *Pak J Biol Sci*. 2008;11:92-7.
26. Lai QX, Bao ZY, Zhu ZJ, Qian QQ, Mao BZ. Effects of osmotic stress on antioxidant enzymes activities in leaf discs of PSAG12- IPT Modified gerbera. *J Zhejiang Univ Sci B*. 2007;8:458-64.
27. Liu ZJ, Guo YK, Bai JG. Exogenous hydrogen peroxide changes antioxidant enzyme activity and protects ultrastructure in leaves of two cucumber ecotypes under osmotic stress. *J Plant Growth Regul*. 2010;29:171-83.
28. Della Croce C, Bronzetti G, Cini M, Caltavuturo L, Poi G. Protective effect of lipoic acid against hydrogen peroxide in yeast cells. *Toxicol in Vitro*. 2003;17: 753-9.
29. Moini H, Packer L, Saris NEL. Antioxidant and prooxidant activities of α -lipoic acid and dihydrolipoic acid. *Toxicol Appl Pharm*. 2002;182:84-90.
30. Ksouri R, Megdiche W, Debez A, Falleh H, Grignon C, Abdely C. Salinity effects on polyphenol content and antioxidant activities in leaves of the halophyte *Cakile maritime*. *Plant Physiol Biochem*. 2007;45:244-9.
31. Foyer CH, Noctor G. Ascorbate and glutathione: the heart of the redox hub. *Plant Physiol*. 2011;155:2-18.