

CAPSELLA RUBELLA TGA4, A BZIP TRANSCRIPTION FACTOR, CAUSES DELAYED FLOWERING IN ARABIDOPSIS THALIANA

Maofu Li^{1,2,3}, Hua Wang^{1,2,3}, Yuan Yang^{1,2,3} and Wanmei Jin^{1,2,3,*}

¹ Institute of Forestry and Pomology, Beijing Academy of Agriculture and Forestry Science, Beijing, 100093, China

² Beijing Engineering Research Center for Deciduous Fruit Trees, Beijing, 100093, China

³ Key Laboratory of Biology and Genetic Improvement of Horticultural Crops (North China), Ministry of Agriculture, Beijing, 100093, China

*Corresponding author: jwm0809@163.com

Received: April 14, 2015; Revised: July 14, 2015; Accepted: July 16, 2015; Published online: December 22, 2015

Abstract: Flowering time is usually regulated by many environmental factors and endogenous signals. TGA family members are bZIP transcription factors that bind to the octopine synthase element, which has been closely linked to defense/stress responses. Most TGA factors interact with *non-expressor of PR1 (NPR1)* and plant defense responses are strengthened by this interaction. *TGA1* and *TGA4* factors bind to *NPR1* only in salicylic acid (SA)-induced leaves, suggesting that *TGA4* has another function during plant development. Here, we isolated a bZIP transcription factor gene, *TGA4*, from *Capsella rubella*. *TGA4* transcripts were detected in most tissues, with high expression in leaves, low expression in stems and flowering buds, and undetectable in siliques. *CruTGA4* was overexpressed in *Arabidopsis thaliana* wild type Col-0 plants. Flowering time and total leaf number in the transgenic plants showed that overexpression of *CruTGA4* could delay flowering in *A. thaliana*. Our findings suggest that *TGA4* may act as flowering regulator that controls plant flowering.

Key words: A bZIP transcript factor; *TGA4* gene; delayed flowering

Abbreviations: bZIP – basic leucine zipper; NPR1 – non-expressor of PR1; qRT-PCR – quantitative real-time polymerase chain reaction; FMI – floral meristem identity; SA – salicylic acid; EMSA – electrophoretic mobility shift assays; CO – constans; FT – flowering locus T; ocs – octopine synthase; GST – glutathione S-transferase; CaMV – cauliflower mosaic virus; MS – Murashige and Skoog; ORF – open reading frame; bp – base pair; NJ – neighbor-joining; LD – long day; SOC1 – suppressor of overexpression of CO 1; WT – wild type

INTRODUCTION

Floral initiation during the transition from vegetative to reproductive growth is one of the key developmental processes in flowering plants. Many studies have indicated that the integration of multiple environmental factors and endogenous signals regulates this transition. Genetic analysis of the long-day annual plant *Arabidopsis thaliana* has identified five major flowering time regulatory pathways (photoperiod, temperature and vernalization, gibberellin, autonomous and aging pathways) that control floral initiation (Srikanth and Schmid, 2011). More and more floral-pathway in-

tegrator genes downstream of the floral pathways are being isolated (Blazquez and Weigel 2000; Hepworth et al., 2002; Kardailsky et al., 1999; Kobayashi et al., 1999; Lee et al., 2000; Moon et al., 2003; Nilsson et al., 1998; Samach et al., 2000). These floral pathway integrator genes cause the floral transition by activating a set of genes known as floral meristem identity (FMI) genes to promote floral development (Blazquez and Weigel 2000; Hepworth et al., 2002; Kardailsky et al., 1999; Lee et al., 2000; Moon et al., 2003; Samach et al., 2000). The photoperiod pathway channels inputs from light, day length, and the circadian clock to promote the floral transition. The B-box transcription factor

CONSTANS (*CO*) is a central regulator of this pathway and *CO* can trigger the production of the mobile florigen hormone *FLOWERING LOCUS T* (*FT*) that induces flower differentiation (Valverde 2011).

TGA family members, a group of bZIP transcription factors, can bind to the octopine synthase (*ocs*) element, which is a plant promoter sequence believed to be involved in defense/stress responses (Foley and Singh 2004). *Ocs* elements were found to regulate the expression of a number of plant pathogen-responsive genes. In *Arabidopsis*, *ocs* element-like sequences are important for the expression of specific glutathione S-transferase (*GST*) and pathogenesis-related genes (Chen and Singh 1999, Lebel et al., 1998). Most TGA factors interact with *NPR1* in *Arabidopsis* (Despres et al., 2003, Kim et al., 2008, Zhang et al., 1999). However, the *TGA1* and *TGA4* factors do not interact with *NPR1* in *Arabidopsis*, only binding to *NPR1* in SA-induced leaves (Despres et al., 2003); this suggests that *TGA4* factors have another function. *TGA4* can bind to the promoter of the *FT* gene according to electrophoretic mobility shift assays (EMSA) and can form a protein complex by interacting with *CO* in the yeast two hybrid system (Song et al., 2008). This suggests that *TGA4* may have some function in controlling plant flowering, but there is no evidence of such activity *in vivo*.

Capsella rubella, an *Arabidopsis* relative, originated through an extreme population bottleneck. This was very important in understanding the common mechanism of adaptation in *Capsella rubella* compared to the model plant *Arabidopsis thaliana*. To obtain insight into the possible function of *TGA4* in plant development, we isolated a *TGA4* gene from *Capsella rubella*. An overexpression vector, fused to the cauliflower mosaic virus (CaMV) 35S promoter, was constructed and used to transform into *Arabidopsis* wild-type Columbia-0 (Col-0) by the floral dip method. Flowering time and total leaf number showed that overexpression of *CruTGA4* could delay flowering in *Arabidopsis*. Our findings suggest that *TGA4* acts as a flowering regulator.

MATERIALS AND METHODS

Plant growth conditions

Arabidopsis thaliana seeds of ecotype Col-0 wild type and *Capsella rubella* were surface-sterilized with 75% alcohol for 2 min and washed at least five times with sterile water, dispersed on 1/2 Murashige and Skoog medium supplemented with 0.8% agar and 2% sucrose, stratified at 4°C for 3 days in the dark, and then germinated at 21°C. The plants were grown in plastic pots containing a mixture of substrate and vermiculite (3:1). Controlled environmental conditions were provided in growth chambers (21°C and 50% relative humidity). The plants were illuminated with fluorescent lights (100-150 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Long day (LD) conditions consisted of 16 h of light followed by 8 h of darkness. The plants were fertilized with mineral nutrient solution (Estelle and Somerville, 1987).

Genomic DNA extraction, PCR amplification and sequencing

Total genomic DNA was extracted from 250 mg of fresh leaf with an EZ Spin Column Genomic DNA Isolation Kit (Biomega Inc., Foster City, CA, USA). Pairs of primers (Supplemental Table 1) were synthesized by Shanghai Sangon (Shanghai, China). The PCR reaction mix contained 0.5 μM of each primer, 25 ng template including genomic DNA and cDNA (see below), 2.5 mM of each deoxynucleotide triphosphate, 2.5 units of KOD Hot Start polymerase (Toyobo Life Science, Osaka, Japan), 1 \times PCR buffer and 2.5 mM MgSO_4 . The PCR program included an initial step of 94°C for 3 min, 35 cycles of 94°C for 15 s, 55°C for 30 s, 68°C for 1-2 min, and a final extension at 68°C for 8 min. The PCR products were separated on 1% agarose gels, stained with ethidium bromide (EtBr), and visualized with a gel-imaging system. The PCR fragments were purified using an EZ-10 Spin Column DNA gel extraction kit (Bio Basic Inc., Markham, Ontario, Canada). PCR products were ligated into the pEASY-T1 vector (TransGen Biotech, China), cloned into the *Escherichia coli* strain DH5 α and the positive clones were sequenced.

Sequence alignment and phylogenetic analyses

Full-length amino acid sequence alignments of TGA4 proteins were performed using ClustalW (Chenna et al., 2003). The genetically mobile structural domains of TGA proteins were determined and annotated using SMART (<http://smart.embl-heidelberg.de/>). Phylogenetic analysis of the *TGA4* genes was based on deduced amino acid sequences. A phylogenetic tree was generated with MEGA version 5.0 using the neighbor-joining (NJ) method with 1000 bootstrap replicates (Tamura et al., 2011).

Expression pattern analysis by qRT-PCR

To investigate the spatial and temporal expression patterns of *TGA4*, qRT-PCR experiments were performed on various organs. Total RNA was extracted from each tissue and first-strand cDNA was synthesized with a RevertAid First Strand cDNA synthesis kit (Thermo Scientific, Waltham, MA, USA) according to the manufacturer's instructions. We performed qRT-PCR with SYBR® Premix Ex Taq™ II (Bio-Rad, Hercules, CA, USA) on a Bio-Rad 480 Real-Time PCR Detection System. Primers used for qRT-PCR are shown in Table S1. Two independent biological replicates and three technical replicates for each sample were performed by qRT-PCR. Data were analyzed using the $2^{-\Delta\Delta CT}$ method as outlined by Livak and Schmittgen (2001). The expression levels of specific genes were normalized to *tubulin*.

Vector construction

To confirm the *in vivo* function of the *TGA4* gene, we created overexpression constructs by PCR cloning. A genomic DNA fragment from *C. rubella* containing the entire coding region, the upstream intergenic region and the downstream intergenic region was amplified by PCR using KOD Hot Start polymerase with gene-specific primer sets (Table S1). The PCR condition was the same as for cloning the *TGA4* gene. The PCR products were purified with a High Pure PCR purification kit (Bio Basic Inc., Markham, Ontario, Canada). The PCR products and the vector pCAMBIA1300 were digested with *Kpn* I and *Spe* I endonucleases. The di-

gested PCR and vector products were ligated with T4 DNA ligase enzyme (Takara Biotechnology, Dalian, China), the positive clones were sequenced and the right clone was used in transgenic tests.

Transgenic tests and phenotypic data

The resulting binary vector was introduced into *Agrobacterium* strain GV3101 and used to transform the *Arabidopsis thaliana* Col-0 wild-type by the floral dip method (Clough and Bent 1998). Transformed seeds were selected on half-strength MS plates containing 50 mg/L hygromycin. Transgenic plants were grown in plastic pots containing a mixture of substrate and vermiculite (3:1) and identified by PCR using the primer of the hygromycin gene (Table S1).

The total leaf number of transgenic plants and the Col-0 wild-type were scored as the number of leaves in the rosette (excluding cotyledons) plus the number of leaves in the inflorescence at the time of opening of the first flower; flowering time was scored by counting the number of days from sowing to the opening of the first flower. Statistical data were analyzed by SPSS 11.5 (SPSS Inc., Chicago, IL, USA) and Sigmaplot 10.0 (Systat Software Inc., San Jose, CA, USA).

RESULTS

Analysis of TGA4 in *Capsella*

TGA4 was amplified from leaf cDNA and the PCR product was cloned and sequenced in pEASY-T1. The full-length sequence of *TGA4* was 1095 bp and contained an open reading frame (ORF) of 364 amino acid residues (Fig. S1A). To study the structure of the *TGA4* gene, we cloned the genomic sequence of *TGA4* from gDNA with the same primers that were used for cDNA cloning (Fig. S1B and Table S1). We compared the gDNA and cDNA sequences of *CruTGA4* and another six *TGA* genes in *Capsella*. The results revealed that these genes comprised 6-9 exons and 5-8 introns. Exons of 60, 78 and 272 bp in length were much conserved in these genes. Exons of 60 bp were presented in all these genes, exons of 78 bp in all but *CruTGA6*

and exons of 272 bp in all except for *CruTGA7*. However, the introns differed in length. The comparison of exon-intron structures of *TGA4* and the other six *TGA* genes show that these genes were a conserved gene family (Fig. 1A).

Phylogenetic analysis

Sequence SMART (<http://smart.embl-heidelberg.de/>) and plant transcription factor database (<http://planttfdb.cbi.pku.edu.cn/>) searches revealed that *CruTGA4* (Carubv10001361m.g) is located on *C. rubella* chromosome 6, that it is 1095 bp long, and that it encodes a 364 amino acid protein with a predicted molecular mass of 41.6 kDa. Amino acids 76-136 contain a typical bZIP basic region domain, which may serve as an active domain. All TGA proteins are characterized by an extraordinarily long ZIP domain consisting of multi-heptad repeats (Fig. 1B and Fig. 2A).

To analyze the phylogenetic relationships between members of the *TGA4* gene family, we performed a phylogenetic analysis of *CruTGA4* and *TGA4* genes from other angiosperms. An NJ phylogenetic tree based on deduced amino acid sequences was generated. In the tree, *CruTGA4* formed a cluster with *CgrTGA4*, *AthTGA4*, *BstTGA4*, *AarTGA4*, *AlyTGA4*, *BnaTGA4* and *EsaTGA4*; together they formed a monophyletic clade in Brassicaceae (Fig. 2B). Therefore, these results further indicate that *TGA4* proteins, as a type of bZIP transcription factor, are conserved not only in Brassicaceae but also in other angiosperms.

Expression patterns of *TGA4*

To understand the functions of *TGA4* gene in *C. rubella* development, we studied the organ-specific expression of *TGA4* using total RNAs derived from different plant tissues and organs by qRT-PCR analysis. Results showed that *TGA4* transcripts were present in most tissues tested, with high expression in leaves, low expression in stems and flowering buds, and undetectable expression in siliques (Fig. 3). This indicated that *TGA4* was expressed during plant development, which is consistent with previous reports (Song et al., 2008).




Fig. 1. Genomic organization and bZIP protein domains of TGA family members in *C. rubella*. **A** – Genomic organization of TGA family members in *C. rubella*. Boxes represent exons and lines represent introns. Numbers indicate the lengths of exons and introns in base pairs. **B** – bZIP protein domains of TGA family members including *CruTGA4* and other six *TGA* genes.

Over-expression of *TGA4* cause delayed flowering in *Arabidopsis*

To assess the potential role of the *CruTGA4* gene in the control of flowering-time, the gene was overexpressed in *Arabidopsis*. A DNA clone of the *TGA4* gene was generated from genomic DNA of *C. rubella* plants, fused to the cauliflower mosaic virus (CaMV) 35S promoter (35S:*CruTGA4*) and introduced into *Arabidopsis* WT Col-0 plants by the floral dip method (Fig. S1B-D). Fifteen transgenic 35S:*TGA4* plants were obtained from half-strength MS plates containing 50 mg/L hygromycin and grown in plastic pots containing a mixture of substrate and vermiculite. The result of PCR using hygromycin gene primers showed that these 15 plants were transgenic lines (Fig. S2 and Table S1). Compared with non-transgenic plants, in the transgenic lines' the flowering time was longer. In the non-transgenic line, the flowering time was about 38




Fig. 2. Phylogenetic relationships among plant TGA4. **A** – Alignment of the deduced amino acid sequences of TGA4s. The bZIP basic region domain is underlined. **B** – An unrooted NJ phylogenetic tree of CruTGA4 and other TGA4 proteins. Bootstrap values from 1000 replicates are shown at each node. TGA4 sequences were obtained from *Malus domestica* (MdoTGA4), *Phoenix dactylifera* (PdaTGA4), *Zea mays* (ZmaTGA4), *Aegilops tauschii* (AtaTGA4), *Triticum urartu* (TurTGA4), *Cucumis sativus* (CsaTGA4), *Vitis vinifera* (VviTGA4), *Medicago truncatula* (MtrTGA4), *Arabidopsis thaliana* (AthTGA4), *Arabidopsis lyrata* (AlyTGA4), *Arabidopsis arenosa* (AarTGA4), *Brassica napus* (BnaTGA4), *Capsella rubella* (CruTGA4), *Oryza sativa* (OsaTGA4), *Capsella grandiflora* (CgrTGA4), *Sorghum bicolor* (SbiTGA4), *Panicum virgatum* (PavTGA4), *Brachypodium distachyon* (BdiTGA4), *Mimulus guttatus* (MguTGA4), *Solanum lycopersicum* (SlyTGA4), *Eucalyptus grandis* (EgaTGA4), *Populus trichocarpa* (PtrTGA4), *Linum usitatissimum* (LusTGA4), *Manihot esculenta* (MesTGA4), *Eutrema salsugineum* (EsaTGA4), *Boechera stricta* (BstTGA4), *Glycine max* (GmaTGA4), *Fragaria vesca* subsp. *vesca* (FveTGA4), and *Prunus persica* (PpeTGA4). Protein sequence information for these genes is given in Table S2.




Fig. 3. Expression analysis of *CruTGA4* in different organs of *C. rubella*. FL – flowering bud; LF – leaf; ST – stem; SL – silique. The relative expression values for *CruTGA4* are relative to *tubulin* and represent means of two biological replicates. SD – standard deviation.

days on average; in plants overexpressing the gene, the flowering time was about 58 days (Fig. 4A). Statistical analysis showed that the transgenic lines had delayed flowering compared with WT Col-0 plants (Fig. 4B).

DISCUSSION

In this study, we isolated a *TGA4* gene from *C. rubella*. *CruTGA4* had a conserved domain typical of bZIP transcription factors; the bZIP motif plays an

important role in determining the function in *Arabidopsis*. The genomic organization of the TGA family is well conserved in Brassicaceae. In the phylogenetic tree, *CruTGA4* was most closely related to TGA4 orthologs from Brassicaceae. Our sequence alignment and phylogenetic tree analysis suggest that *CruTGA4* should have a similar function to other *TGA4* orthologs. It was reported that TGA family transcription factors, which interact with *NPR1*, are closely linked to defense/stress responses (Zhou et al., 2000, Zhang et al., 1999). However, *TGA4* factors do not interact with *NPR1*, and bind to *NPR1* only in SA-induced leaves (Despres et al., 2003, Kim et al., 2008, Zhang et al., 1999). Thus, *TGA4* factors might have multiple roles in plant development and may have other functions besides regulating defense/stress and nitrate responses.

Analysis of *TGA4* expression patterns showed that *TGA4* transcripts were present in most tissues tested, and were highly expressed in leaves (Fig. 3); this result is in accordance with Song et al. (2008). *CO* has been demonstrated to promote flowering by activating the expression of *FT* and *SUPPRESSOR OF OVEREXPRESSION OF CO 1 (SOC1)* (Kardailsky et al., 1999, Kobayashi et al., 1999, Samach et al., 2000). The diurnal expression patterns of *TGA4* were similar to those of *CO* during the night in both LD and short day conditions (Song et al., 2008). So we infer that *TGA4* may interact with *CO* and bind to the *FT* promoter, and the




Fig. 4. Overexpression of *CruTGA4* delays flowering in *Arabidopsis* under long day conditions. **A** – Representative images of flowering wild type and transgenic overexpression plants grown under LD conditions (16/8 h, light/dark). Photography at 52 days after sowing. **B** – Flowering time and number of rosette leaves. *** significant at $P < 0.001$ compared with wild-type levels.

TGA4 mRNA abundance mirrors the diurnal oscillations of *CO*. Mutations in *FT* delay flowering in wild-type and in 35S:*CO* plants (Koornneef et al., 1991; Onouchi et al., 2000). *FT* mRNA accumulates in the night under LD conditions (Suarez-Lopez et al., 2001), so *TGA4* may regulate the expression of *FT* through its interaction with *CO* during the dark period, and thereby control plant flowering. Our transgenic plants showed that overexpression of the *CruTGA4* gene delayed flowering in *A. thaliana* in long day conditions. This finding suggests that overexpression of *CruTGA4* may repress the expression of *FT* through its interaction with *CO*, and then delay flowering in *A. thaliana*. Although the flowering time between wild-type and transgenic plants demonstrated that *CruTGA4* is involved in the regulation of plant flowering, further genetic studies using mutants of *co*, *ft* and *tga4* genes, are needed for functional complementation experiments and analysis of the relationships among *TGA4*, *CO* and *FT* using transgenic lines.

Acknowledgments: This work was supported by the National Science Foundation for Young Scholars of China (31201606), the Important Crops Molecular Breeding Technology Innovation (KJ CX20140202), Special Fund for the Construction of Scientific and Technological Innovation Capability (No. KJ CX20140110), and Beijing Municipal Science and Technology Commission (No. D131100000113001) projects.

Authors' contributions: Maofu Li and Wanmei Jin conceived and designed the experiments; Maofu Li and Hua Wang performed the experiments; Maofu Li and Yuan Yang analyzed the data, Maofu Li and Wanmei Jin wrote the paper.

Conflict of interest disclosure: The authors declare no conflict of interest.

REFERENCES

- Blazquez, M.A. and D. Weigel (2000). Integration of floral inductive signals in *Arabidopsis*. *Nature*. **404**, 889-892.
- Chen, W. and K.B. Singh (1999). The auxin hydrogen peroxide and salicylic acid induced expression of the *Arabidopsis* *GST6* promoter is mediated in part by an ocs element. *Plant J.* **19**, 667-677.
- Chenna, R., Sugawara, H., Koike, T., Lopez, R., Gibson, T.J., Higgins, D.G. and J.D. Thompson (2003). Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res.* **31**, 3497-3500.
- Clough, S.J. and A.F. Bent (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**, 735-743.
- Despres, C., Chubak, C., Rochon, A., Clark, R., Bethune, T., Desveaux, D. and P.R. Fobert (2003). The *Arabidopsis* NPR1 disease resistance protein is a novel cofactor that confers redox regulation of DNA binding activity to the basic domain/leucine zipper transcription factor *TGA1*. *Plant Cell.* **15**, 2181-2191.
- Estelle, M.A. and C. Somerville (1987). Auxin-resistant mutants of *Arabidopsis thaliana* with an altered morphology. *Mol. Gen. Genet.* **206**, 6.
- F. Valverde (2011). CONSTANS and the evolutionary origin of photoperiodic timing of flowering. *J. Exp. Bot.* **62**, 2453-2463.
- Foley, R.C. and K.B. Singh (2004). *TGA5* acts as a positive and *TGA4* acts as a negative regulator of ocs element activity in *Arabidopsis* roots in response to defence signals. *FEBS Letters.* **563**, 141-145.
- Hepworth, S.R., Valverde, F., Ravenscroft, D., Mouradov, A. and G. Coupland (2002). Antagonistic regulation of flowering-time gene *SOC1* by *CONSTANS* and *FLC* via separate promoter motifs. *EMBO J.* **21**, 4327-4337.
- Kardailsky, I., Shukla, V.K., Ahn, J.H., Dagenais, N., Christensen, S.K., Nguyen, J.T., Chory, J., Harrison, M.J. and D. Weigel (1999). Activation tagging of the floral inducer *FT*. *Science.* **286**, 1962-1965.
- Kim, M.G., Kim, S.Y., Kim, W.Y., Mackey, D. and S.Y. Lee (2008). Responses of *Arabidopsis thaliana* to challenge by *Pseudomonas syringae*. *Mol. Cells.* **25**, 323-331.
- Kobayashi, Y., Kaya, H., Goto, K., Iwabuchi, M. and T. Araki (1999). A pair of related genes with antagonistic roles in mediating flowering signals. *Science.* **286**, 1960-1962.
- Koornneef, M., Hanhart, C.J. and J.H. van der Veen (1991). A genetic and physiological analysis of late flowering mutants in *Arabidopsis thaliana*. *Mol. Gen. Genet.* **229**, 57-66.
- Lebel, E., Heifetz, P., Thorne, L., Uknes, S., Ryals, J. and E. Ward (1998). Functional analysis of regulatory sequences controlling *PR-1* gene expression in *Arabidopsis*. *Plant J.* **16**, 223-233.
- Lee, H., Suh, S.S., Park, E., Cho, E., Ahn, J.H., Kim, S. G., Lee, J.S., Kwon, Y.M. and I. Lee (2000). The AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways in *Arabidopsis*. *Genes Dev.* **14**, 2366-2376.
- Livak, K.J. and T.D. Schmittgen (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* **25**, 402-408.
- Moon, J., Suh, S.S., Lee, H., Choi, K.R., Hong, C.B., Paek, N.C., Kim, S.G. and I. Lee (2003). The *SOC1* MADS-box gene integrates vernalization and gibberellin signals for flowering in *Arabidopsis*. *Plant J.* **35**, 613-623.
- Nilsson, O., Lee, I., Blazquez, M.A. and D. Weigel (1998). Flowering-time genes modulate the response to *LEAFY* activity. *Genetics.* **150**, 403-410.

- Onouchi, H., Igeno, M.I., Perilleux, C., Graves, K. and G. Coupland (2000). Mutagenesis of plants overexpressing CONSTANS demonstrates novel interactions among *Arabidopsis* flowering-time genes. *Plant Cell*. **12**, 885-900.
- Samach, A., Onouchi, H., Gold, S.E., Ditta, G.S., Schwarz-Sommer, Z., Yanofsky, M.F and G. Coupland (2000). Distinct roles of CONSTANS target genes in reproductive development of *Arabidopsis*. *Science*. **288**, 1613-1616.
- Song, Y.H., Song, N.Y., Shin, S.Y., Kim, H.J., Yun, D.J., Lim, C.O., Lee, S.Y. Kang, K.Y. and J.C. Hong (2008). Isolation of CONSTANS as a TGA4/OBF4 interacting protein. *Mol. Cells*. **25**, 559-565.
- Srikanth, A. and M. Schmid (2011). Regulation of flowering time: all roads lead to Rome. *Cell Mol. Life Sci*. **68**, 2013-2037.
- Suarez-Lopez, P., Wheatley, K., Robson, F., Onouchi, H., Valverde, F and G. Coupland (2001). CONSTANS mediates between the circadian clock and the control of flowering in *Arabidopsis*. *Nature*. **410**, 1116-1120.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and S. Kumar (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731-2739.
- Zhang, Y., Fan, W., Kinkema, M., Li, X. and X. Dong (1999). Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid induction of the *PR-1* gene. *Proc. Nat. Acad. Sci. USA*. **96**, 6523-6528.
- Zhou, J.M., Trifa, Y., Silva, H., Pontier, D., Lam, E., Shah, J. and D.F. Klessig (2000). *NPR1* differentially interacts with mem-

bers of the TGA/OBF family of transcription factors that bind an element of the *PR-1* gene required for induction by salicylic acid. *Mol. Plant Microbe Interact*. **13**, 191-202.

Supplementary Material (available online)

Supplementary Fig. S1. The cloned *CruTGA4* gene and overexpression vector construct. The *CruTGA4* gene cloned from cDNA (A) and gDNA (B). (C) A diagram of the *CruTGA4* overexpression vector. (D) Confirmation of the overexpression vector construct by double enzyme digestion, 1-9 indicate the different positive clone, CK – empty vector.

Available at: <http://serbiosoc.org.rs/sup/SupplMat/FIGS1.tif>

Supplementary Fig. S2. Confirmation of transgenic plants by PCR using the hygromycin gene primer. The numbers 1-15 indicate the 15 transgenic plants, 'p' represents the plasmid of the overexpression vector construct, 'WT' indicates the non-transgenic plant.

Available at: <http://serbiosoc.org.rs/sup/SupplMat/FIGS2.tif>

Supplementary Table S1. Primers used in this study.

Available at: <http://serbiosoc.org.rs/sup/SupplMat/SupplTable1.doc>

Supplementary Table S2. Sequence information of proteins mentioned in Fig. 2B.

Available at: <http://serbiosoc.org.rs/sup/SupplMat/SupplTable2.doc>